

RICHARD J. NEVYS
onstantinov A. S. 1962 - Ves nekotorych vodnykh bezpozvonochnykh kak funktsia ich lineinykh zmerov - Nauč. Dokl. vyssh. Šk. 3: 17-20.
onstantinov A. S. 1969 - Ves životnykh kak funktsia ich lineinykh rozmerov - Ž. obšč. Biol. 3: 16-272.

uferov V. P. 1956 - Nekotorye dannye o chiščonom pitanii ličinek *Tendipedidae* - Dokl. Akad. Nauk SSSR, 111: 466-470.
uferov V. P. 1961 - O pitanii ličinek *Pelopitinae* (Diptera, *Tendipedidae*) - Trudy Inst. Biol. i odochran. 4: 232-245.

awacz W. 1969 - The characteristics of sinking materials and the formation of bottom deposits in entrophic lake - Mitt. int. Verein. Limnol. 17: 319-331.
orduchaj B. 1954 - Materialy po sravneniu vesu vodnykh bezpozvonochnykh ssena Dona - Trudy probl. temat. Sovetsk. zool. Inst. 2: 223-241.
andie I. H. 1959 - The diurnal activity of the layer invertebrates at the surface of Lac La Ronge, Saskatchewan - Can. J. Zool. 37: 945-956.

ezynski E., Kajak Z. 1965 - Investigations on the mobility of the bottom fauna in the lakes Białe, Mikolajskie and Śniardwy - Bull. Acad. pol. Sci. Cl. II. Sér. Sci. Biol. 13: 345-353.
odubnaja T. L., Sorokin I. I. 1961 - Głubina sloja optymalnogo pitanija tubificid v svjazi s perebneženijem v grunte - Byull. Inst. Biol. i odochran. 10: 14-17.
Horacka J. 1963 - Phytoplankton in lakes near Wegorzewo in the light of peculiarities of the environment - Pol. Arch. Hydrobiol. 11: 189-217.

opincev B. A. 1949 - O skorosti razloženia organičeskogo veščstva otmeršego planktona - Izv. vses. gidrobiol. Obšč. 1: 34-43.
odniewska I. 1967 - Dynamics of the abundance and biomass of phytoplankton in Lakes sołajskie and Tańkowskie - Ecol. pol. A, 15: 139-153.
odniewska I. 1974 - The structure and production of phytoplankton in Mikolajskie lake - Ecol. pol. 22: 66-105.

ančykowska A. 1961 - Gwałtowna redukcja liczebności *Dreissena polymorpha* Pall. w kilku jeziorach mazurskich okolic Mikolajek - Ecol. pol. B, 7: 151-153.
ančykowska A. 1964 - On the relationship between abundance, aggregations and "condition" *Dreissena polymorpha* Pall. in 36 Mazurian Lakes - Ecol. pol. A, 12: 635-690.
ančykowska A., Przytcka-Jusiak M. 1968 - Variations in abundance and biomass of microbenthos in three mazurian lakes - Ecol. pol. A, 16: 539-559.

umiec J. 1963 - Wpływ roślinności wynurzonej i sposobu jej koszenia na faunę denną stawów - Acta hydrobiol. 5: 315-335.
al J. M. 1957 - Community metabolism in a temperate cold spring - Ecol. Monogr. 27: 233-301.
kers T. F. 1969 - The turnover ratio in production ecology of freshwater invertebrates - Am. J. Zool. 103: 173-185.
jczak-Migała I. 1965 - Wpływ dwuletniego zalania stawów na faunę denną - Roczn. Nauk i Techn. 1: 215-227.
ker W. 1961 - Lebenszyklus und Vertikalverteilung der *Chironomidae* (Dipt.) *Sergentia corecina* T. und Titisee - Verh. int. Verein. Limnol. 14: 962-967.

Paper prepared by J. Stachowiak

AS' ADDRESSES:
hab. Zdzisław Kujek
ysztof Durąg
hydrobiologii
Ekologii PAN
ów Lesny k. Warszawy
Łomianki

Krzysztof LEWANDOWSKI and Anna STAŃCZYKOWSKA

Department of Hydrobiology, Institute of Ecology, Polish Academy of Sciences, Dziekanów Leśny near W.

THE OCCURRENCE AND ROLE OF BIVALVES OF THE FAMILY UNIONIDAE IN MIKOŁAJSKIE LAKE*

ABSTRACT: The distribution, numbers, biomass, annual production and age structure of *Unionidae* were investigated. A mathematical relation between the size (age) and weight of a bivalve was determined for two species: *Anodonta piscinolis* Nilsson and *Unio tumidus* Philipsson. The filtration abilities and the role of these invertebrates in Mikolajskie Lake were determined.

Contents

- 1. Introduction
2. Area and methods
3. Results
3.1. Species composition of Unionidae; distribution in the lake
3.2. Age structure
3.3. Growth and the relation of weight and size
3.4. Biomass and annual production
3.5. Filtration possibilities; the role of Unionidae in the lake
3.6. Final remarks
4. Summary
5. Polish summary (Streszczenie)
6. References

I. INTRODUCTION

The considerable attention given to the family Unionidae (Żadin 1938, Agrell 194 Ökland 1963, Petrov 1964, Negus 1966, Widuto and Kompowski 196 Piechocki 1969, Tudorancea 1972 and others) is because its members belong the largest freshwater invertebrates and the contribution of bivalves to benthic biomass sometimes 99% (Ökland 1963) and thus they are of considerable significance in the biocenosis of a water body. The members of the family Unionidae belong to the large

*Praca wykonana w ramach problemu węzłowego nr 09.1.7 (grupa tematyczna "Procesy decydujące o czystości powierzchniowych wód śródlądowych").

freshwater filtering organisms. These animals purify the water from suspension and excrete faeces which are abundant in organic matter and thus are a suitable life environment for several benthic organisms. The moving *Unionidae* stir the bottom deposits and the accumulation at the bottom of shells of dead individuals which decompose with difficulty cause the increasing shallowness of water bodies.

The aim of this paper was to study the distribution, numbers, annual production and age structure of bivalves of the family *Unionidae* in Mikołajskie Lake. The daily filtration of single *Unionidae* and their numbers allowed to estimate the filtration possibilities for the whole lake.


Detailed study on these animals in Mikołajskie Lake has not been carried out before. There is only a remark about four species of the family *Unionidae* in Mikołajskie Lake given by Berger (1966) who describes the malacofauna of Masurian Lakeland.

2. AREA AND METHODS

Mikołajskie Lake is a typical water body in Great Masurian Lakeland (Fig. 1). This is a tunnel-valley lake, eutrophic, holomictic, of a surface 460 ha, maximum depth 27.8 m, mean depth 11.0 m. The littoral occupies 19% of the total lake surface (Kajak, Hillbricht-Ilkowska and Pieczyńska 1972). The shores of the lake on the southern part are usually high and steep, up to 30 m above the water level. The shallow part of the littoral is narrow and rapidly gains depth (Synowiec 1961). Emergent vegetation covers 39.2 ha, of which 32.3 ha overgrown with reed — *Phragmites communis* Trin. (Kowalcwski and Wasilewski 1966). The reed grows on most of the littoral zone of Mikołajskie Lake, but it does not grow at all in the north-eastern part where is the town Mikołajki. In places not covered with reed, at small depths, the bottom is stony or sandy but is always covered by a thin layer of mud.

The *Unionidae* were collected an 10 sites evenly distributed along Mikołajskie Lake (Fig. 1). The sites were chosen according to the variety of littoral types — distribution of reed and various species of submerged plants, configuration and type of bottom and changes in the environment as a result of man's activities (sewage). On each site, several metres broad and running along the shore, samples were taken starting at the depth of 0.5 m and then at 1.5, 2.5, 3.5 m and etc. down to the depth at which the *Unionidae* practically do not occur. At shallow places (0.5 m) the bivalves were picked by hand on a framed surface ($0.5 \times 0.5 \text{ m} = 0.25 \text{ m}^2$) chosen at random at the bottom. This was repeated 10 times on each site. At deeper places the material was collected using the bottom dredge of a side 0.4 m and drawn along the bottom in parallel to the shore from a boat at anchor. The dredge was drawn 5 times at each depth in particular sites. Thus obtained material was used to estimate the numbers of *Unionidae* in the lake. Of the two commonly applied methods of quantitative catches of bivalves (among others of Widuto and Kompowski 1968, Stańczykowska and Nowicki 1966) the frame method is undoubtedly more precise. The species of bivalves were identified as well as their age and they were measured and weighed fresh and dry (after drying up to constant weight). In order to observe the changes in the distribution of *Unionidae* in the lake during the season the material was collected four times between May and September 1972. The fluctuations of the water level in the lake during the studies did not exceed 10 cm.

In August the studies on the intensity of water filtration by *Unionidae* were also carried out using the method for *Dreissena polymorpha* (Pallas), in Mikołajskie Lake (Mattice, Stańczykowska and Ławacz 1972, Stańczykowska, Ławacz and



tice - in press). A specially constructed, on the principle of a flow respirometre, experimental set was used to estimate the filtration intensity, feeding rate, assimilation and respiration in organisms examined. One experimental set consisted of 6 cages; in 4 cages *Unio*, or *Anodonta* was placed and 2 cages without bivalves were the control. The set was lowered to the lake on a rope to the depth of 3 m.

3. RESULTS

Species composition of *Unionidae*; distribution in the lake

Quantitative studies carried out in Mikołajskie Lake have shown the occurrence of *Unionidae* species in brackets - the share in total numbers: *Anodonta piscinalis* Nilsson (4.0%), *Anodonta cygnea* L. (4.0%), *Anodonta (Pseudanodonta) complanata* Rossmäslér (20.5%), *Unio tumidus* Philipsson (20.5%), *Unio pictorum* L. (3.8%).

In the whole 477 individuals of *Unionidae* were caught. *Anodonta piscinalis* dominated in the lake and *Unio tumidus* was quite frequently caught. Other species occurred sporadically. The same species, with the exception of *Anodonta (Pseudanodonta) complanata*, are mentioned by Berger (1960) for Mikołajskie Lake. In this study this species has been mentioned as occasionally rare.

Similar species composition is recorded in several lakes of the Masurian Lakeland (own data, Berger 1960, Widuto and Kompowski 1966), in water bodies of other parts of Poland (Klimowicz 1958, Zaćwilichowska 1965, Soszka 1968) and in lakes of many European regions (Žadin 1938, Agrell 1949, Ökland 1963, Todorancea 1972).

In Mikołajskie Lake the *Unionidae* appear at the depth of about 0.2 m. At the depth of 0.5 m the numbers of *Unionidae* were the largest (0.76 ind./m², fresh weight with shell 4.0 g/m², the shell 4.0 g, dry weight without shell 0.49 g/m²). Together with the increasing

Tab. 1. Mean abundance and biomass of *Unionidae* in different depth of Mikołajskie Lake

A - abundance (ind./m²), B - wet biomass with shell (g/m²), C - shell weight (g/m²), D - dry biomass without shell (g/m²)

Depth (m)	<i>Anodonta piscinalis</i>				<i>Unio tumidus</i>				<i>Unionidae</i> - total			
	A	B	C	D	A	B	C	D	A	B	C	D
0.5	0.33	7.4	1.7	0.26	0.31	4.5	1.90	0.19	0.76	16.2	4.0	0.49
1.5	0.43	8.7	2.1	0.35	0.09	0.9	0.30	0.03	0.60	12.2	3.4	0.44
2.5	0.42	7.4	1.6	0.22	0.10	1.0	0.30	0.03	0.53	9.4	2.3	0.24
3.5	0.27	4.4	1.0	0.16	0.07	0.2	0.07	0.02	0.43	5.9	1.5	0.19
4.5	0.15	2.5	0.5	0.08	0.02	0.1	0.03	0.01	0.19	3.1	0.8	0.09
5.5	0.05	1.1	0.2	0.03	0	0	0	0	0.05	1.1	0.2	0.03

The determination of fresh weight of bivalves is always difficult due to the presence of the shell which contains mainly inorganic substances and the water content in the mantle cavity (troublesome to remove in small animals). Although the water evaporates after the animal's death the water in tissues also dries up, and especially in the case of young age classes. Thus the results cannot be without an error.

depth the numbers of *Unionidae* decrease evenly (Fig. 2, Tab. 1). This study has covered the zone down to 5.5 m deep not only because the *Unionidae* occur sporadically there but at the depth of 4.5 m. The most common in Mikołajskie Lake *Anodonta piscinalis* occurred in large numbers at the depth of 1.5-2.5 m. The majority, over 50% of all specimens of species *Unio tumidus*, were collected at 0.5 m. At greater depths this species was found in very small numbers and its range of occurrence ended at the depth of 4.5 m. Of other species *Anodonta cygnea* occurred at depths of 1.5-4.5 m and dominated at shallow parts, whereas *Unio pictorum* occurred at 0.5-3.5 m and dominated in deeper parts.




Fig. 2. Percentage of *Unionidae* at particular depths in Mikołajskie Lake

Similar data on the distribution of *Unionidae* in other lakes are given by several authors. *Unionidae* usually occur in the littoral zone, e.g. in Kortowskie Lake (Masurian District) to a depth of 4 m (Widuto and Kompowski 1966), in lake Borrevann (Norway) to

6 m deep (Ökländ 1963). They dominate at the depth of 0.5 m, e.g. in Kortowskie Lake (Widuto and Kompowski 1968), in lakes Velikoe and Svyatoe in the USSR (Zadin 1938). In Kortowskie Lake *Unio tumidus* dominates at the depth of 0.5 m in Widuto and Kompowski (1968) and in lake Crapina (the Danube Delta) at the depth of 0.4 m (Tudorancea 1972) — similarly as in Mikotajskie Lake. But as regards the *Anodonta piscinalis*, it is most abundant at greater depths in other lakes as well — at 2 m in Kortowskie Lake (Widuto and Kompowski 1968), at 1.5 m in lake Družno (Zimek 1960), at 2–3 m in lake Zurich (Burla 1972), at 3 m in lake Borrevain (Ökländ 1963). Thus *Unio tumidus* dominates at the depth of about 0.5 m and *Anodonta piscinalis* at about 2 m.

The differences in the density of *Unionidae* on particular sites in Mikotajskie Lake were not great. Despite considerable environmental differences the densities were similar and low (mean numbers — 0.4 ind./m², fresh weight with shell — 9.9 g/m², the shell 2.4 g, dry weight without shell 0.3 g/m²) on the majority of sites. The exception is site 3 where the density was three times higher as compared to the mean one. This site, at the depth of 0.5 m, had the largest number of *Unionidae* in June (7.2 ind./m², fresh weight with shell — 154.1 g/m², the shell — 2 g, dry weight without shell — 4.61 g/m²). The numbers of *Unio tumidus* were also the highest there (6.4 ind./m²). *Anodonta piscinalis* did not occur in such large numbers and were evenly distributed. Site 3 had a sandy bottom covered with a thin mud layer and considerable amount of plant remains, the reed did not grow abundantly there. The shore was steep with plenty of trees and bushes the branches of which hung over the water. Apart from the very steep shore site 3 did not differ much from sites 4 and 9. However, the numbers of *Unionidae* were much lower on these sites.

Although the numbers on particular sites do not differ much the *Unionidae* were less numerous on stony bottom than on a sandy-mud or mud bottom. But these differences were as great as in other water bodies (among others Zadin 1938, Widuto and Kompowski 1968). It seems that aquatic vegetation does not affect much the distribution of *Unionidae* and this is also confirmed by the data from the literature (Widuto and Kompowski 1968). In Mikotajskie Lake the densely growing reed have affected the movements of these animals. On site 10 a dense reed belt divided the vast littoral bed without *Unionidae* from the open water. The *Unionidae* occurred in the place where the reed belt was lower and less dense.

Site 7 was at the outlet of municipal sewage. The water there (acc. to data of Pieczyńska, Sikorska and Ozimek 1975) contained considerable amounts of ammonia nitrogen (up to 20 mg N-NH₄/l) and also of chlorides, Na⁺, K⁺, Ca²⁺, its oxygen saturation was high thus proving a considerable inflow of organic substances accompanied by ger deficits. The strong wave action in Mikotajskie Lake results in quick dilution of sewage concentrations of particular substances are already observed at the distance of 5 m from sewage outlet (Pieczyńska, Sikorska and Ozimek 1975). On site 7 the *Unionidae* did not occur in the direct vicinity of sewage inflow but 15–20 m farther, the numbers there were close to the mean values. Pedras (unpublished data) observed similar situations in the occurrence of *Anodonta* sp. in polluted environments of Mikotajskie Lake. In Kortowskie Lake the numbers of *Unionidae* were smaller in the region of municipal sewage outlet (Widuto and Kompowski 1968).

Over the summer months the density of *Unionidae* did not change much on particular sites with large and small numbers.

But there are some regularities in the distribution of *Unionidae* in the summer and at various depths (Fig. 3). The distribution of *Anodonta piscinalis* does not change much although numbers increase considerably at 0.5 m in August and September. The occurrence of *Anodonta piscinalis* is much more even than of the second species as regards the abundance *Unio tumidus* — *Unio tumidus*, in June and August prefers the depth of 0.5 m, whereas in May and September — greater depths.




Fig. 3. Percentage of *Unionidae* at particular depths in Mikotajskie Lake in the summer 1972

3.2. Age structure

The members of various age classes (different shell length) of the *Unionidae* are found at different depths (Fig. 4). At the depth of 0.5 m the large individuals of *Anodonta piscinalis* and *Unio tumidus* are usually found. The depth of 1.5 m is colonized by all classes of age of species *Anodonta piscinalis*. Smaller individuals of this species (10–30 mm long) occur abundantly at depths of 2.5–3.5 m. The smallest specimens of *Unio tumidus* are also the most abundant at greater depths (2.5–4.5 m). Similar results have been obtained by Widuto and Kowalski (1968) in Kortowskie Lake. This has been slightly different in lake Borrevann




Fig. 4. Percentage of *Unionidae* with different shell lengths at particular depths in Mikotajskie Lake (summer 1972)

(Ökland 1963). Young *Anodonta piscinalis* dominated both in shallow places (0.4 m in deep (6 m), older individuals were caught at 3–5 m.

The age of caught braves has been determined using the method of counting growth on shells. This method (among others) is discussed in detail by Crowley (1957) is comm applied to specify the age of *Unionidae* (among others Ökland 1963, Negus 1969, Magini and Stanezowska 1971). The oldest specimen *Anodonta piscinalis* caught in Mikotajskie Lake have been 9–10 years old. The age structure




Fig. 5. Age structure of *Unionidae* in Mikotajskie Lake (1972): *Anodonta piscinalis* (A) 303 individuals, *tumidus* (B) 87 individuals

lected *Anodonta piscinalis* and *Unio tumidus* is presented in Figure 5. Individuals 4–6 years old are the majority. It is understandable that the number of older individuals is smaller due to natural mortality. But the small number of the youngest specimens is quite striking. This has been pointed out by several authors studying the *Unionidae* in various water bodies and streams among others Ökland 1963, Petrov 1964, Negus 1966, Piechocki 1969, Tudorancea 1969, 1972, Magnin and Stańczykowska 1971). The scientists frequently come across the young specimens in rivers where the *Unionidae* form beds (Petrov 1964, Piechocki 1969). The beds in the river Grabia (observations by Piechocki 1969) and in the river Oka (Petrov 1964) have been formed mainly by adult individuals, and there were very few young.

The difficulties in finding young *Unionidae* may be due to their relatively small size and can be caused by applying a method appropriate for large benthic organisms or caused by the toxic biology of these animals. The type of reproduction and the occurrence of the larval stage (glochidium) which is a parasite of fish may be responsible for the considerable spreading, whereas the adults tend to concentrate as a result of their sexual maturity (Piechocki 1969) and they mature rather late: in the case of *Anodonta* at the age of 2–4 years and in the case of *Unio* at the age of 2–3 years (among others Žadin 1938, Piechocki 1969). It there can be a different hypothesis, such as the occurrence of greater aggregations of young *Unionidae* in places frequently visited by fish — carriers of glochidia — and it is just a sheer coincidence that the majority of scientists do not find them. According to Piechocki's (pers. comm.) observations there are such aggregations in the river Grabia. Negus (1966) has observed this type of aggregations of young *Unionidae* in the Thames and the transition from a static to free way of living of the *Unionidae* coincides with the reproduction activity of fish. This is the problem of the occurrence of young *Unionidae* is not yet explained and requires some other studies.

It should be indicated that the age structure of a given species is considerably affected by seasonal fluctuations of numbers of individuals of the new generation and this depends on several factors. Sometimes the *Unionidae* do not reproduce at all at a certain year as it happened in 1965 in lake Jijila (Tudorancea 1972).

3.3. Growth and the relation of weight and size

A characteristic feature of the *Unionidae* is their intensive growth in the first years and afterwards less intensive but continuing all their life. The character of growth of *Anodonta piscinalis* and *Unio tumidus*, despite the differences in size, is similar (Fig. 6). The limits of the length range of successive age classes form a curve which is typical for the growth of *Unionidae* — individuals having such measurements have been undoubtedly under best conditions. The curve of mean shell lengths for particular classes differs considerably from the former although it slightly shows a quicker increment of shell length in earlier years and its rate later on.

Between the length of a bivalve (age) and its weight there is a relation:

$$Y = a X^b$$

where Y — weight, X — length of a bivalve, a and b — parameters. According to empirical data




Fig. 6. Mean length and the range of length of *Unionidae* depending on age (Nikolajskie Lake 1972)




Fig. 7. Weight of a bivalve as a function of shell length (Nikolajskie Lake 1972)
1 — fresh weight (with shell), 2 — dry shell weight, 3 — dry body weight (without shell)

ne values of parameters a and b were estimated for the relation of total weight (fresh weight with shell), weight of the shell itself and dry weight to the length of both species *Anodonta piscinalis* and *Unio tumidus*. Estimations of these parameters allowed to find model types of relations² (Fig. 7). Similar relations for the *Unionidae* have been also found by Oklinda (1963) and Widuto and K o m p o w s k i (1968). The increase of length at a young age is usually accompanied by a small weight increment, whereas at older age when the shell length does not change much the weight of *Unionidae* rapidly increases. As the exponent (b) is an increase of weight unit in relation to an increase of length unit thus the increase in weight for determined shell length is greater the greater the exponent. In the case of *Anodonta piscinalis* the shell weight increases proportionally the quickest as regards the three weights examined.

Tab. II. Shell weight, dry biomass and content of water of *Anodonta piscinalis* in dependence according to age

Age (years)	Shell weight (%)	Dry biomass without shell (%)	Water (%)
1	4.8	0.9	94.3
2	15.3	1.6	88.1
3	17.9	2.0	80.1
4	23.5	3.1	73.4
5	25.5	2.9	71.6
6	26.7	2.9	70.4
7	26.6	2.8	70.6
8	25.5	2.7	71.7

Tab. III. Shell weight, dry biomass and content of water of *Unio tumidus* in dependence according to age

Age (years)	Shell weight (%)	Dry biomass without shell (%)	Water (%)
1	26.0	1.2	72.8
2	25.6	2.2	72.2
3	33.3	2.2	64.5
4	35.0	2.9	62.2
5	41.5	2.5	56.0
6	42.7	3.0	54.3
7	42.2	3.0	54.8
8	42.8	4.8	52.4

whereas in the case of *Unio tumidus* the dry body weight. The total weight increase slowest for both species (Fig. 7). This is connected with the decreasing amount of water animal's body together with the age and corresponds to the percentage increase of content of the shell and dry body to the total weight of a bivalve (Tabs. II and III). These individuals in both species but the general regularities are similar.

3.4. Biomass and annual production

In the estimates of the total number of individuals and biomass of *Unionidae* in Mikolajskie Lake the age structure of the dominant species (*Anodonta piscinalis*) has been used. It is assumed that the number estimations are not standard for the youngest specimens (Fig. 7) therefore the number of 1, 2, 3 and 4 year olds in the lake is at least the same as the number of 5 year olds. Thus the assumed number of *Unionidae* was 680,000 individuals (biomass 12.5 tons fresh weight with shells, this including 3 tons of shells, dry weight without shells 210 kg). These numbers are for an area of 114 ha which is the littoral zone of the lake down to 5.5 m deep (Synowicz 1961).

The calculations of annual production of *Unionidae* in Mikolajskie Lake were made using the method applied by Greze (1965), Kajak and Rybak (1966) and other methods of study annual production of various aquatic organisms. This method is used by many authors studying bivalves (Negus 1966, Tudorancea and Florescu 1968a, 1968b; Magnin and Stanczykowska 1971). Table IV presents particular stages of calculating the annual production by *Anodonta piscinalis*. Particular age classes have their numbers (n), mean weight without shell (b) and mean annual biomass increase of an individual (Δb) and according to these data the biomass, annual production, and the ratio of production to biomass (P/B) are calculated. The biomass of total *Anodonta piscinalis* population in Mikolajskie Lake is 54.9 kg/ha and the annual production 18.2 kg/ha. The ratio P/B has a mean value of 0.33 but changes together with the age from higher values for younger to lower values for older classes of age (Tab. IV). For all *Unionidae* in Mikolajskie Lake the biomass is 65.6 kg/ha and the annual production 22.4 kg/ha and annual P/B 0.34. The small number of members of species *Anodonta cyganea* makes it impossible to calculate precisely its biomass and annual production. However, the approximate estimations show that the ratio P/B is very high -- 0.66. It is lower for *Unio tumidus* ($P/B = 0.28$).

The biomass of *Unionidae* in Mikolajskie Lake is relatively small. For example, in Crapina (the Danube Delta) the biomass of *Anodonta piscinalis* is 161.7 kg/ha and the annual production 50.9 kg/ha (Tudorancea and Florescu 1969). The corresponding values for *Unio tumidus* are 116.9 kg/ha and 29.9 kg/ha (Tudorancea and Florescu 1968b). The ratio P/B is very similar in Mikolajskie Lake and in lake Crapina. In the latter P/B for *Anodonta piscinalis* 0.32 on average, 0.76 for sexually immature individuals and 0.17 for older classes of age (Tudorancea and Florescu 1969). Mean annual P/B for *Unio tumidus* is 0.25 and is also higher for young classes of age (0.71) and lower for older classes (0.17) (Tudorancea and Florescu 1968b). The biomass given by Negus (1966) for the Thames is even higher: biomass of *Anodonta anatina* L. (= *A. piscinalis*) 648.2 kg/ha, biomass of all *Unionidae* 1207.3 kg/ha. But the P/B values are much lower the comparison to the corresponding data in Mikolajskie Lake. P/B for *Anodonta anatina* 0.20 and for *Unio tumidus* 0.12, for all *Unionidae* 0.16. These low P/B values indicate that in these populations older individuals prevail. Nevertheless, it should be indicated that the assumed number

²The authors wish to acknowledge Dr. T. Wierzbowska for these estimates carried out in the Laboratory of Statistics and Modelling, Institute of Ecology, Polish Academy of Sciences.

Tab. IV. Biomass and annual production of *Anodonta piscinalis* in Mikotajskie Lake (1972)

Age (years)	Numbers of individuals per 1 m ² <i>n</i>	Biomass of one individual (wet weight without shell) (g) <i>b</i>	Increase of the biomass of one individual per year (g) Δb	Average biomass (g/m ²) $B = n \cdot b$	Production per year (g/m ²) $P = n \cdot \Delta b$	<i>P/B</i>
1	0.06*	1.2	1.2	0.07	0.07	1.00
2	0.06*	3.2	2.0	0.19	0.12	0.62
3	0.06*	8.1	4.9	0.49	0.29	0.58
4	0.06*	14.4	6.3	0.86	0.38	0.43
5	0.06	19.2	4.8	1.15	0.29	0.25
6	0.05	23.3	4.1	1.16	0.21	0.18
7	0.03	33.0	9.7	0.99	0.29	0.29
8	0.005	59.3	17.3	0.25	0.09	0.35
9	0.005	66.0	15.7	0.33	0.08	0.24
Total	0.390	—	—	5.49	1.82	0.33

*Theoretical numbers (= numbers of the 5th class of age).

ing in Mikotajskie Lake is higher than it has been found (this has been already mentioned). If calculations are made on the basis of the existing age structure then the *P/B* index will be also lower due to smaller contribution of young (e.g. it would be 0.29 for *Anodonta piscinalis*).

3.5. Filtration possibilities; the role of *Unionidae* in the lake

Partly on the basis of the results of experiments and partly on the basis of the literature data filtration possibilities of *Unionidae* in Mikotajskie Lake were estimated. In August, two field experiments lasting 24 hours were carried out to study filtration, consumption and assimilation of food in *Unionidae* belonging to the middle class of age (4–5 years). The set of animals was the same in both experiments, i.e. two *Anodonta piscinalis* individuals with shells 60 and 54 mm long, and two *Unio tunidus* individuals with shells 48 and 54 mm long. Every few hours measurements turbidity of water that passed through experimental and control cages was measured. Thus during the experiment the filtration of animals examined could be observed; periods of decreased filtration intensity were irregular, either shorter or longer. The percentage of used up available seston ranged from 10 to 80% (47% on average). The daily feeding rate was 20.6 mg dry seston weight per individual on average. The *Unionidae* excreted 4.2 mg faeces/ind. per 24 hr on average. Assimilation of food calculated from the equation $A = C - F$ (— assimilation, *C* — consumption, *F* — faeces) was on average 16.4 mg/ind. per 24 hr (79.6% in proportion to the filtrated seston). Both species had similar filtration rates — they filtered on average 300 ml water per hour (= 14.3 ml/hr per g fresh weight with shell) but at individual variability ranging from 60 to 490 ml/hr. Similar values were obtained by Skresenskiy and Kondratev, cited by Alimov (1965), and according to them *Anodonta*

piscinalis and *Unio tunidus*, 46 mm long, filtrate 400 ml water per hour. The filtration rate of *Anodonta cygnea*, 80–100 mm long is 250–300 ml/hr according to De Bruin and Davids (1970).

For the calculations of filtration possibilities of *Unionidae* populations in Mikotajskie Lake including all classes of age the equation given by Alimov (1965) was used:

$$V = 84.14P^{-0.51}$$

for the relation between filtration rate — *V* (ml/hr per g) and weight of *Anodonta piscinalis* (*P* (g)). The filtration rates obtained for the field experiments in Mikotajskie Lake are similar to analogous values calculated according to the equation above. Although in these calculations only *Anodonta piscinalis* (most abundant in the lake) is taken into account the results related to the total number of *Unionidae* in the lake as the filtration possibilities of other species belonging to this family are similar. Table V presents the results of this equation for par-

Tab. V. Filtration of *Anodonta piscinalis* in various age classes

Age (years)	Average biomass of individual (g) <i>P</i>	Filtration* (ml/hr per 1 g of animal body) <i>V</i>	Filtration (ml/hr per animal) <i>V · P</i>
1	1.25	76.5	96
2	3.80	44.3	168
3	9.90	27.1	268
4	18.80	19.5	366
5	25.80	16.5	425
6	31.80	15.0	477
7	44.90	12.6	566
8	65.00	10.4	676
9	88.00	8.9	783

*According to Alimov (1969).

classes of age of *Anodonta piscinalis*. According to these data the *Unionidae* in Mikotajskie Lake can filtrate during 24 hours about 5000 m³ of water and during the whole vegetation season (i.e. about 6 months) the thirtieth part of all epilimnion down to 7 m-deep a filtrate from water about 2.5 tons dry seston weight.

3.6. Final remarks

Generally it can be said that in Mikotajskie Lake there are the same *Unionidae* species as in other water bodies. The domination relations are also similar — usually the most abundant species are *Anodonta piscinalis* and *Unio tunidus*. Still the numbers of *Unionidae* are much smaller in other water bodies and the same can be said of biomass and annual production. These numbers are much higher in Kortowskie Lake where the mean value in the zone of occurrence is 10.6 ind./m² and at the inflow of ground waters rich in humic substances the ma-

number of 256 ind./m² is found (Widuto and Kompowski 1968). The numbers of the order 200-400 individuals/m² are given by Zadin (1928) for rivers with dams and lakes. A similar number has been recorded (300 ind./m²) in the dam reservoir Goczakowice Krzyżanek, pers. comm.). According to Agrell (1949) the abundance of *Unionidae* depends on the extent of lake trophy. And thus he gives the number of 1.8 individuals of *Anodonta piscinalis* per 1 m² of oligotrophic and mesotrophic lakes in southern Sweden, 2 ind./m² of eutrophic lakes and 20-30 ind./m² of highly eutrophic lakes. Together with the increasing trophy the numbers of *Unio pictorum* increase and of *Unio tumidus* decrease. In similar lakes of the same geographical regions some authors observe extremely different numbers of *Unionidae*, e.g. in lakes Julia and Crapina (the Danube Delta) 3.8 and 11.5 ind./m² Tudorancea (1972), in lakes Saint-Louis and Deux Montagnes (south-eastern Canada) 1 and 25 ind./m² (Magnin and Stachykowski 1971) and in lakes Parovoye and Elkhoe (central European part of the USSR) 4 and 65 ind./m² at the depth of 0.5 m Zadin (1938). In water bodies where *Unionidae* are very abundant their percentage in total benthic biomass is very high. In lake Borrevam in which *Anodonta piscinalis* is the only species of the family *Unionidae* its mean number is 10 ind./m² and as regards its weight 95.7% of total benthic fauna in this lake (Öklund 1963). In Mikołajskie Lake the number of *Unionidae* is small as well as their percentage in the biomass of entire bottom fauna although molluscs dominate in this lake and the contribution of *Dreissena polymorpha* to benthic biomass is largest in its zone of occurrence (0.2-8.0 m) and has the mean number of 300 ind./m² Stachykowski et al. - in press). The effect of bivalves on the lake biocenosis is primarily connected with their filtration. All populations of bivalves in Mikołajskie Lake (bivalves of the family *Unionidae*, *Dreissena polymorpha*, bivalves of the family *Sphaeriidae*) in filtrate during the vegetation season a double amount of water contained by the epilimnion of this lake. The dominant role is that of *Dreissena polymorpha* (Stachykowski et al. - in press).

The majority of material used here is from the master's thesis of Mr. K. Lewandowski "Selected problems in the ecology of bivalves of the family *Unionidae* in Mikołajskie Lake" which has been carried out in the Department of Hydrobiology, Zoological Institute, University of Warsaw, under the guidance of Asst. Prof. Dr. Ewa Peczyńska. The work was given the award of Polish Hydrobiological Society in 1974.

4. SUMMARY

The study on bivalves of the family *Unionidae* was carried out in eutrophic Mikołajskie Lake (Masurian Lakeland) (surface area 460 ha, mean depth 11.6 m) on 10 sites (Fig. 1) and the material was collected four times between May and September 1972. Five *Unionidae* species were found. *Anodonta piscinalis* Nilsson (1.5%) and *Unio tumidus* Philipsson (20.5%) were the most abundant species. Other species: *Anodonta* sp. L., *Unio pictorum* L. and *Anodonta (Pseudanodonta) complanata* Kossmanster were found sporadically. The numbers of *Unionidae* in the littoral zone down to 5.5 m deep compared to the data of other authors were relatively low (mean value 0.4 ind./m², maximum value 7.2 ind./m²). The largest numbers were served at small depths: at 1.5-2.5 m *A. piscinalis*, at 0.5 m *U. tumidus* (Fig. 2) and during the summer season there were some changes in the distribution of various species at particular depths which were very ar in the case of *U. tumidus* (Fig. 3). The most frequently caught *Unionidae* were those 4-6 years old, and individuals were rarely found (Fig. 5). And thus in further estimates (biomass, production and filtration of *Unionidae* in this lake) higher numbers of younger classes of age were used than those found in light of the shell only and the dry body weight was determined (Fig. 7). Annual production of *Unionidae* Mikołajskie Lake was 22.4 kg/ha at a biomass 65.6 kg/ha (annual means of $P/B = 0.34$). The mean annual for *A. piscinalis* was 0.33 (Tab. IV), 0.28 for *U. tumidus* and 0.66 for *A. cygnea*. Data on the filtration of *Unionidae* were obtained in field experiments. Four and five year old *A. piscinalis* and *U. tumidus*

(49-60 mm long) filtrated about 300 ml per hour. Their daily feeding rate was on average 20.6 mg (season weight per individual, and faeces production - 4.2 mg. Food assimilation in the summer seas 79.6% in relation to filtrated seston. According to the data presented and the literature data on the fill of *Unionidae* it was estimated that over the period of one vegetation season all *Unionidae* in Mikołajskie (680,000 individuals) can filtrate the thirtieth part of the total volume of epilimnion in this lake a filtrate from water about 2.5 tons dry seston weight.

5. POLISH SUMMARY (STRESZCZENIE)

Badania nad małzami z rodziny *Unionidae* prowadzono w eutroficznym jeziorze Mikołajskim 460 ha, gł. śr. 11,6 m), wyznaczając na nim 10 stanowisk (fig. 1) i zbierając materiał czterokrotnie w maju - wrześniu 1972 r. Stwierdzono występowanie pięciu gatunków *Unionidae*. Najpospolitsi: *Anodonta piscinalis* Nilsson (71,5%) i *Unio tumidus* Philipsson (20,5%). Pozostałe sporadycznie spotykano: *Anodonta cygnea* L., *Unio pictorum* L. i *Anodonta (Pseudanodonta) complanata* Kossan. Liczebność *Unionidae* w strefie litoralnej do głębokości 5,5 m była, w porównaniu z danymi innymi w stosunkowo niską (średnio - 0,4 osobnika/m², maksimum - 7,2 osobnika/m²). Największe liczebności stwierdzane były na małych głębokościach: na głębokości 1,5-2,5 m - *A. piscinalis*, na głę 0,5 m - *U. tumidus* (fig. 2), przy czym w ciągu sezonu letniego notowane były pewne zmiany w liczbie różnych gatunków na poszczególnych głębokościach, szczególnie wyraźnie u *U. tumidus* (Najliczniej potrawiano *Unionidae* w średnim wieku, mające 4-6 lat. Stosunkowo rzadko były znalezione osobniki młode (fig. 5). Toteż do dalszych szacunkowych obliczeń (biomasy, produkcji i filtracji *Unionidae* w tym jeziorze) przyjęto wyższą liczebność młodszych klas wiekowych niż stwierdzono w rzeczywistości. Dla dwóch gatunków *Unionidae* ustalono zależność między długością małża (wiekiem) ciężarem całkowitym, ciężarem samej muszli i ciężarem suchego ciała (fig. 7). Produkcja roczna *Unionidae* jeziora Mikołajskiego wynosi 22,4 kg/ha przy biomacie - 65,6 kg/ha (średnie roczne $P/B = 0,34$). U *A. piscinalis* średnie roczne P/B wynosi 0,33 (tab. IV), u *U. tumidus* - 0,28 a u *A. cygnea* - 0,66. W rymentach terenowych uzyskano dane o filtracji *Unionidae*. *A. piscinalis* i *U. tumidus* w wieku 4-5 lat (średnio 48-60 mm) filtrują około 300 ml wody na godzinę. Ich dobowa racja pokarmowa wynosi 20,6 mg suchej masy sestonu na osobnika, fekalia - 4,2 mg. Asymilacja pokarmu w okresie letnim - 79,6% w stosunku do odfiltrowanego sestonu. Opierając się na własnych i literaturowych danych o *Unionidae* oszacowano, że w ciągu jednego sezonu wegetacyjnego wszystkie *Unionidae* występujące w jeziorze, odfiltrować zaś mogą z wody około 2,5 tony suchej masy sestonu.

6. REFERENCES

1. Agrell J. 1949 - The shell morphology of some Swedish Unionides as affected by ecological factors - Ark. Zool. 41 A, 15: 1-39.
2. Alimov A. F. 1965 - Filtrationnaja sposobnost' moljuskov roda *Sphaerium* (Scopoli) - Dokl. Naik. SSSR, 164: 195-197.
3. Alimov A. F. 1969 - Nekotorye obščie zakononomnosti processa filtracii u dvustvorčatykh i skov - Ž. obšč. Biol. 30: 621-631.
4. Berger L. 1969 - Badania nad małzami (*Mollusca*) Pojezierza Mazurskiego - Kadan. fizjog. zashed. 6: 7-49.
5. Burla H. 1972 - Die Abundanz von *Anodonta*, *Unio pictorum*, *Unio pictorum*, *Unio pictorum* auris und *Lymnaea ovata* im Zürichsee, in Abhängigkeit von der Wassertiefe und zu verschiedenen J. eiten - Vjschr. Naturforsch. Ges., Zürich, 117: 129-151.
6. Crowley T. E. 1957 - Age determination in *Anodonta* - J. Couch. 24: 201-207.
7. De Bruin J. P. C., David C. 1970 - Observations on the rate of water pumping of the water mussel *Anodonta cygnea zelandica* (Gmelin) - Neth. J. Zool. 20: 380-391.
8. Grece V. N. 1965 - Temp rosta i produktivnye vozmozhosti populacii ryb - Cidrobiol. Ž. 1.
9. Kajak Z., Hillbrecht-Ilkowska A., Pieczyńska E. 1972 - The production of *Unionidae* in several Polish lakes (In: Productivity problems of freshwaters, Eds. Z. Kajak, A. Hillbrecht-Ilkowska - PWN, Polish Scientific Publishers, Warszawa-Kraków, 129-147).
10. Kajak Z., Rybak J. J. 1966 - Production and some trophic dependences in benthos

primary production and zooplankton production of several Masurian lakes — Verh. Internat. Verein. Limnol. 16: 441—451.

1. K i m e k L. 1960 — Studia nad fauną denną jeziora Drużno — Zesz. nauk. Univ. Mikołaja Kopernika Toruń, 7: 29—59.

2. K l i m o w i c z H. 1958 — Mączaki Zalewu Wiślanego i zależności ich rozmieszczenia od zasolenia — Pol. Arch. Hydrobiol. 5: 93—123.

3. K o w a l c z e w s k i A., W a s i l e w s k i L. 1966 — Differentiation of biomass of *Phragmites communis* Trin. and its production in Mikołajskie Lake — Bull. Acad. pol. Sci. Cl. II, 2: 219—223.

4. M a g n i n E., S t a Ń c z y k o w s k a A. 1971 — Quelques données sur la croissance, la biomasse et la production annuelle de trois mollusques *Unionidae* de la région de Montréal — Can. J. Zool. 49: 491—497.

5. M a t t i c e J., S t a Ń c z y k o w s k a A., Ł a w a c z W. 1972 — Feeding and assimilation of *Dreissena polymorpha* in Mikołajskie Lake, Poland — Am. Zool. 12: 209—209.

6. N e g u s C. L. 1966 — A quantitative study of growth and production of unionid mussels in the river Thames at Reading — J. Anim. Ecol. 35: 513—532.

7. Ō k l a n d J. 1963 — Notes on population density, age distribution, growth and habitat of *Anodonta piscinalis* Nils. (Moll. Lamellibr.) in a eutrophic norwegian lake — Nytt. Mag. Zool. 11: 19—43.

8. P e t r o v V. V. 1964 — K ocenke promysłowej močności populacji moljuskow semejstva *Unionidae* — Izv. Inst. ozern. rečn. rybn. Choz. 57: 42—46.

9. P i e c h o c k i A. 1969 — Obserwacje biologiczne nad małżami z rodziny *Unionidae* w rzecze Grabzi — Acta Hydrobiol. 11: 56—67.

10. P i e c z y n s k a E., S i k o r s k a U., O z i m e k T. 1975 — The influence of domestic sewage on the littoral zone of lakes — Pol. Arch. Hydrobiol.

11. S o s z k a G. 1968 — Selected problems of the ecology of molluscs (*Mollusca*) of the brackish lake Żebisko — Ekol. pol. Ser. A, 16: 729—733.

12. S t a Ń c z y k o w s k a A., Ł a w a c z W., M a t t i c e J. (in press) — Use of field measurements of filtration and assimilation in evaluating *Dreissena polymorpha* in lake ecosystem — Pol. Arch. Hydrobiol.

13. S t a Ń c z y k o w s k a A., Ł a w a c z W., M a t t i c e J., L e w a n d o w s k i K. (in press) — Bivalves as a factor effecting circulation of matter in Lake Mikołajskie — Limnologia.

14. S t a Ń c z y k o w s k a A., N o w i c k i M. 1966 — Próba zastosowania metody swobodnego nurkowania w badaniach nad małżem — *Dreissena polymorpha* Pall. — Ekol. pol. Ser. B, 12: 331—335.

15. S y n o w i e c A. 1961 — Morfologia Jeziora Mikołajskiego — Dokument. geogr. PAN, 2: 52—69.

16. T u d o r a n c e a C. 1969 — Comparison of the populations of *Unio tumidus* Philipsson from the complex of Crapina-Jiřilla marshes — Ekol. pol. Ser. A, 11: 185—204.

17. T u d o r a n c e a C. 1972 — Studies on *Unionidae* populations from the Crapina-Jiřilla complex of peeds (Danube zone liable to inundation) — Hydrobiologia, 39: 527—561.

18. T u d o r a n c e a C., F l o r e s c u M. 1968a — Cu privire la fluxul energetic al populatiei de *Unio pictorum* din Balta Crapina — An. Univ. Bucuresti Ser. Stiint. Natur. 17: 233—243.

19. T u d o r a n c e a C., F l o r e s c u M. 1968b — Considerations concerning the production and energetics of *Unio tumidus* Philipsson population from the Crapina marsh. — Trav. Mus. Hist. nat. Gr. Antipa, 8: 395—409.

20. T u d o r a n c e a C., F l o r e s c u M. 1969 — Aspecte de productiei si energietici populatiei de *Anodonta piscinalis* Nilsson din Balta Crapina — Studii Cerc. Biol. 21: 43—55.

21. W i d u t o J., K o m p o w s k i A. 1968 — Badania nad ekologią małżów z rodziny *Unionidae* w jeziorze Korzkoskim — Zesz. nauk. wyższ. Szk. roln. Olszt. 24: 479—497.

22. Z a ć w i l i c h o w s k a K. 1965 — Benthos in littoral of the Goczalkowice Reservoir in 1958—1959 — Acta Hydrobiol. 7: 83—97.

23. Ż a d i n V. I. 1938 — Semejstwo *Unionidae* — Fauna SSSR, Moljuskii, nov. ser. 18: 1—169.

Paper prepared by J. I. Rybak

FOR'S ADDRESS:
Grzegorz Lewandowski,
na Stańczykowska,
d Hydrobiologii
zbu Ekologii PAN,
and w Lasny K. Warszawy,

Dormident G. KRYLOV

Department of Biology, State University of Moscow, Moscow

TENDENCY TO GROUPING IN SPATIAL DISTRIBUTION OF SMALL MAMMALS IN A FOREST HABITAT

ABSTRACT: A plot 21.25 ha in the environs of Moscow has been surveyed in April and November 1967, and in April 1968, during 10 days each time. Snap-traps were set a network, 50 x 10 m. In spring and autumn, all the resident population of *Clethrionomys glareolus* (Schreb.) and *Apodemus sylvaticus* (L.) was trapped. It was found to be possible to determine concentration of *Microtus arvalis* (Pall.) in April and July, 1967. A tendency to unite into territorial groupings has been discovered in *C. glareolus* (spring, summer), and *A. sylvaticus* (autumn). Formation of territorial groupings seems to be caused by the low population density of the species, as well as by the existence of interspecific hierarchy in the use of territory.

Contents

1. Introduction
2. Method and material
3. Results
 - 3.1. Intensity of the removal of small mammals
 - 3.2. Spring distribution of animals on the plot
 - 3.3. Discovery of territorial groupings in *Microtus arvalis* in summer
 - 3.4. Autumn distribution of *C. glareolus* and *A. sylvaticus* on the plot
4. Discussion
5. Summary
6. Polish summary (Streszczenie)
7. References

I. INTRODUCTION

The character of spatial distribution of small mammals is one of the more important features of the population. The detailed analysis of the distribution of small mammals in a forest is rather difficult because specific features of the forest given. Nevertheless, the idea of a distribution of small mammals in forest and other plant communities has been accepted years ago. Uneven distribution is conditioned both by the diversity of the habitat and by and interspecific relations among the animals.